

Amendments to the Claims

1. **(Withdrawn-Currently Amended)** A method for the intra-operative treatment of a tumor to inhibit dissemination of tumor cells, which comprises administering to a patient ~~during an intra-operative treatment~~ a preparation consisting of an antibody directed against a tumor-associated antigen and at least one pharmaceutically acceptable carrier selected from the group consisting of ~~an auxiliary substance,~~ a buffer, a salt and a preservative ~~whereby immunocomplexing of tumor cells within the scope of the surgical intervention inhibits dissemination of tumor cells,~~ and wherein the administration of said antibody is carried out ~~within 4 hours prior to surgery and during surgery,~~ and wherein said immunocomplexing of tumor cells activates an antibody-dependent cellular cytotoxicity effector function and a complement dependent cytotoxicity effector function and inhibits tumor cell dissemination.
2. **(Withdrawn)** The method according to claim 1, wherein the antibody is directed against an epitope of a surface antigen of a tumor cell.
3. **(Withdrawn)** The method according to claim 1 or 2, wherein the tumor cells are from an epithelial tumor.
4. **(Withdrawn)** The method according to claim 1, wherein the antibody is directed against an epitope of an antigen selected from the group consisting of peptides, proteins, carbohydrates and glycolipids.
5. **(Withdrawn)** The method according to claim 1, wherein the antibody is in an antibody mixture of various antibodies having a specificity for tumor-associated antigens.
6. **(Canceled)**
7. **(Withdrawn)** The method according to claim 1, wherein the antibody binds to the tumor-associated antigen with an affinity below a Kd value of 10^{-6} mol/l.

8. **(Withdrawn)** The method according to claim 1, wherein the source of said antibody is a mouse or a human.
9. **(Withdrawn)** The method according to claim 1, wherein the antibody is administered systemically in a single dose of at least 50 mg per patient.
10. **(Withdrawn)** The method according to claim 1, wherein the antibody is locally applied to the tumor tissue and/or to the wound area.
11. **(Canceled)**
12. **(Withdrawn)** The method according to claim 1, wherein the surgical intervention is carried out for a biopsy and/or for the removal of a solid tumor.
13. **(Withdrawn)** The method according to claim 1, wherein the surgical intervention is carried out for the purpose of determining the malignancy of a tumor.
14. **(Withdrawn)** The method according to claim 1, wherein immune complexes of the antibody and tumor tissues are determined after the surgical intervention.
15. **(Withdrawn)** The method according to claim 1, wherein immune complexes of the antibody and tumor cells in blood or serum samples are determined.
16. **(Canceled)**
17. **(Withdrawn)** The method according to claim 4, wherein the antigen is a member selected from the group consisting of EpCAM, NCAM, CEA, Lews Y, Sialyl-TN, Globo H, GD2, GD3 and GM2.

18. **(Withdrawn)** The method according to claim 7, wherein said K_d value is 10⁻⁷ mol/l.
19. **(Withdrawn)** The method according to claim 7, wherein said K_d value is 10⁻⁸ mol/l.
20. **(Withdrawn)** The method according to claim 9, wherein said single dose is at least 100 mg.
21. **(Withdrawn)** The method according to claim 9, wherein said single dose is at least 200 mg.
22. **(Withdrawn)** The method according to claim 9, wherein said single dose is at most 2 g.
23. **(Canceled)**
24. **(Canceled)**
25. **(Withdrawn)** The method according to claim 4, wherein said antibody is directed against an epitope of a carbohydrate tumor associated antigen.
26. **(Withdrawn)** The method according to claim 25, wherein said antigen is a member selected from the group consisting of Lewis Y, Glob H, Sialyl-TN, GD2 and GD3.
27. **(Withdrawn)** The method according to claim 26, wherein said antigen is Lewis Y antigen.
28. **(Canceled)**
29. **(Withdrawn-Currently Amended)** A method for the intra-operative treatment of a tumor to inhibit dissemination of tumor cells, which comprises administering to the patient an antibody directed against the tumor-associated antigen Lewis Y ~~during an intra-operative~~

~~treatment whereby immunocomplexing of tumor cells within the scope of the surgical intervention inhibits dissemination of tumor cells~~, and wherein the administration of said antibody is carried out ~~within 4 hours prior to surgery and during surgery~~, and wherein said immunocomplexing of tumor cells activates an antibody-dependent cellular cytotoxicity effector function and a complement dependent cytotoxicity effector function and inhibits tumor cell dissemination.

30. **(Withdrawn)** The method according to claim 29, wherein the antibody is administered during or immediately before the surgical intervention

31. **(Withdrawn)** The method according to claim 29, wherein the antibody is administered during the surgical intervention.

32. **(Withdrawn)** The method according to claim 29, wherein the tumor cells are from an epithelial tumor.

33. **(Canceled)**

34. **(Withdrawn)** The method according to claim 29, wherein the antibody binds to the tumor-associated antigen with an affinity below a K_d value of 10^{-6} mol/l.

35. **(Withdrawn)** The method according to claim 29, wherein said antibody is a human or a mouse antibody.

36. **(Withdrawn)** The method according to claim 29, wherein the antibody is administered systemically in a single dose of at least 50 mg per patient.

37. **(Withdrawn)** The method according to claim 29, wherein the antibody is locally applied to the tumor tissue and/or to the wound area.

38. **(Withdrawn)** The method according to claim 29, wherein the surgical intervention is carried out for a biopsy and/or for the removal of a solid tumor.
39. **(Withdrawn)** The method according to claim 29, wherein the surgical intervention is carried out for a determination regarding the malignancy of a tumor.
40. **(Withdrawn)** The method according to claim 29, wherein immunocomplexes of the antibody and tumor cells in blood or serum samples are determined.
41. **(Withdrawn)** The method according to claim 34, wherein said Kd value is 10^{-7} mol/l.
42. **(Withdrawn)** The method according to claim 34, wherein said Kd value is 10^{-8} mol/l.
43. **(Withdrawn)** The method according to claim 36, wherein said single dose is at most 2 g.
44. **(Canceled)**
45. **(Canceled)**
46. **(Withdrawn)** The method according to claim 1, wherein said antibody is a chimeric antibody or a humanized antibody.
47. **(Withdrawn)** The method according to claim 29, wherein said antibody is a chimeric antibody or a humanized antibody.
48. **(Withdrawn-Currently Amended)** A method for the intra-operative treatment of a tumor to inhibit dissemination of tumor cells, which comprises administering to a patient during an intra-operative treatment a preparation consisting of: i. an antibody directed against a tumor-associated antigen, ii. an adjuvant and iii. and at least one pharmaceutically acceptable carrier selected from the group consisting of an auxiliary substance, a buffer, a salt and a preservative,

~~whereby immunocomplexing of tumor cells within the scope of the surgical intervention inhibits dissemination of tumor cells, and wherein the administration of said antibody is carried out within 4 hours prior to surgery and during surgery, and wherein said immunocomplexing of tumor cells~~ activates an antibody-dependent cellular cytotoxicity effector function and a complement dependent cytotoxicity effector function and inhibits tumor cell dissemination.

49. **(Currently Amended)** A method for the intra-operative inhibition of the dissemination of tumor cells, which comprises administering to a patient during an intra-operative treatment a preparation consisting of an antibody directed against a tumor-associated antigen and at least one pharmaceutically acceptable carrier selected from the group consisting of ~~an auxiliary substance,~~ a buffer, a salt and a preservative ~~whereby immunocomplexing of tumor cells within the scope of the surgical intervention inhibits dissemination of tumor cells, and wherein the administration of said antibody is carried out within 4 hours prior to surgery, during surgery, or both, and wherein said immunocomplexing of tumor cells~~ activates an antibody-dependent cellular cytotoxicity effector function and a complement dependent cytotoxicity effector function and inhibits tumor cell dissemination.